

Computational Models of Memory Formation In Healthy and Diseased Microcircuits of the Hippocampus

Vassilis Cutsuridis¹

¹Institute of Molecular Biology and Biotechnology (IMBB), Foundation for Research and Technology Hellas (FORTH), Heraklion, Crete, Greece

Correspondence:

Vassilis Cutsuridis, Institute of Molecular Biology and Biotechnology (IMBB), Foundation for Research and Technology - Hellas (FORTH), Nikolaou Plastira 100 GR-70013, Heraklion, Crete, Greece. Email: vcutsuridis@gmail.com;

WHAT IS ASSOCIATIVE MEMORY

Memory is important in our lives. Without a memory we are unable to remember our past experiences and our loved ones, yet being able to think about the future. Without memory we cannot learn anything. Associative memory is the ability to learn and remember the relationship between unrelated items, events, and/or objects [68]. When we try to find a specific piece of information in our memory sometimes we do not retrieve it immediately. Our brain, which contains aspects of the present situation and contextual information pointing at the missing piece, starts then a sequential process of associations from one item to the next that eventually leads to the missing piece. Once this piece of information is found, we immediately recognize it as the one we have been searching for.

Associative memory has been one of the oldest artificial neural network (ANN) paradigms. The concept of the associative memory was first introduced by the formalism of a correlation matrix [64, 76, 40, 54, 56, 57], where, memory patterns were encoded as the activity patterns across a network of computing units. Pattern storage was accomplished by Hebbian modification of the connections between the computing units. A memory was recalled when an activity pattern that was a partial or noisy version of a stored pattern was instantiated in the network. Network activity then evolved to the complete stored pattern as appropriate units were recruited to the activity pattern, and noisy units were removed, by threshold-setting of unit activity. Memory capacity for accurate recall was strongly dependent on the form of patterns to be stored and the learning rule employed [57].

An example of an associative memory comprising of 6 input channels interacting with 6 output channels forming a matrix of 36 elements is depicted in Figure 1. Activity in a channel is represented by 1 and inactivity by 0. Associations between the input and output patterns (x^i, y^i), $i = 1, 2, \dots$, are formed via a Hebbian learning rule, where co-activation of the input and output units results in an irreversible transition of synapses from 0 to 1. Such associations between binary stimulus events are then stored in a 6 x 6 connectivity matrix, C . Special cases of associative memory are the auto-association, where $x^i = y^i$ for all pairs i , and the hetero-association, where x^i

$\neq y^i, i = 1, 2, \dots$ [56]. In figure 1A three associations are stored: $x^1 \rightarrow y^1$, $x^2 \rightarrow y^2$, and $x^3 \rightarrow y^3$. The resulting C matrix represents the three sets of paired events.

Recall of an input pattern is accomplished by multiplying the matrix C by a corresponding input pattern, e.g. $X_3 = (1 \ 1 \ 0 \ 1 \ 0 \ 0)$ and performing an integer division analogous to a variable threshold, the value of which is equal to the number of ones in the cueing pattern (i.e. $\theta = 3$). Perfect recall can be achieved by this division process even if the patterns share common active elements, provided that not too many different patterns have been presented (Fig. 1B & C). Errors in recall will begin to occur as the matrix approaches saturation (Fig. 1D).

EARLY VIEWS OF ASSOCIATIVE MEMORY IN HIPPOCAMPUS

David Marr [45, 46] was the first computational scientist to formulate a neural implementation of the correlation matrix in the hippocampus. His network consisted of N principal neurons, one inhibitory neuron and two types of inputs. All neurons were modeled as simple threshold neurons [48] with resting threshold equal to one. Each of the Y inputs strongly depolarized a principal neuron and caused it to fire. All X inputs contacted all principal neurons. Their synaptic weights were initialized to zero and were strengthened according to a Hebbian rule. The X inputs also excited the inhibitory interneurons, which in turn inhibited the principal neuron's somata. The produced inhibitory signal was proportional to the total number of non-zero elements in the input pattern and performed a division operation in the principal neuron's soma allowing this way the neurons that learned the pattern to recall it accurately.

Although Marr's model was a successful one in predicting that the hippocampus works like a content-addressable memory (CAM) system, it was very rudimentary because the types of neurons used in this scheme were simple threshold neuronal nodes and the synaptic weights were updated according to an iterative time quantized update scheme, providing a very rough insight into the dynamical processes in the hippocampus. Since then a dramatic accumulation of knowledge about the morphological, physiological and molecular characteristics, as well as the connectivity and synaptic properties of excitatory and inhibitory neurons in the hippocampus have been

witnessed [14]. Excitatory neurons are primarily pyramidal neurons and they

circuits of associative memory and discuss future challenges.




Figure 1 Example of a correlation matrix (adapted with permission from [12]). (A) Associations of three memory patterns $X^1 \rightarrow Y^1$, $X^2 \rightarrow Y^2$, $X^3 \rightarrow Y^3$ using the mechanism of the correlation matrix. The resulting C matrix represents the three sets of paired events. (B) Perfect recall of an input pattern. (C) Pattern completion of an input pattern. (D) Saturation. Errors in recall start to occur as the matrix approaches saturation.

constitute 90% of all neurons in the hippocampus, whereas the remaining 10% are interneurons, primarily inhibitory, which are classified according to their morphological, physiological, molecular and synaptic characteristics into other numerous subclasses [63]. Collections of thousands of such cells then interact in cell assemblies (microcircuits), with each microcircuit being individual machinery, which receives, processes and transmits information. In the next section, I will briefly review the experimental literature of the hippocampus regarding the different families of neurons and their operations in memory formation and rhythm generation. In section 3, I will review representative examples of simple and detailed spiking neuronal models of associative memory. In section 4, I discuss practical issues and difficulties involved in realistic biophysical modeling of (micro)-

NEURONAL DIVERSITY, MICROCIRCUITS AND RHYTHMS IN THE HIPPOCAMPUS

The hippocampus has been studied extensively yielding a wealth of data on cell types and their passive and active properties, network architecture and synaptic plasticity [14]. Hippocampus contains principal excitatory neurons (granule cells (GCs) in DG and pyramidal cells (PCs) in regions CA3 and CA1) and a large variety of inhibitory interneurons [20, 63]. Neurons in the hippocampus receive external inputs via different pathways from the cortex and the extrahippocampal areas. In the dentate gyrus, GCs receive inputs directly from layer II of the entorhinal cortex (EC). In CA3, inputs come from the EC layer II to the distal apical tree of the PCs. Inputs to proximal and

basal dendrites are largely from other CA3 PCs. Another excitatory input to CA3 comes from the GCs of the dentate gyrus, which form the mossy fiber synapses in the very proximal region of the apical tree of the PCs. In CA1, the Schaffer collateral excitatory input from the CA3 PCs impinges to PC proximal dendritic regions. Recurrent collaterals from other CA1 PCs synapse on the basal dendritic tree, whereas perforant path inputs from EC layer III reach the distal region of the apical dendritic trees of PCs.

Cells in the hippocampal regions compute information differently. DG is implicated in pattern separation [45, 47, 49, 77, 31], CA3 in pattern completion [45, 49] and CA1 in novelty detection [75] and mismatch of expectations [30]. In addition, regions CA3 and CA1 have been proposed to be auto- and hetero-associators for the storage of declarative memories, respectively [69]. Computation in each hippocampal region takes time creating temporal windows of excitability, which are evident by local field potentials (LFPs) [6].

Theta rhythm (4-10 Hz) is one such LFP [1, 72] and it has been shown to play an instrumental role in the coordination of neuronal firing in the entorhinal-hippocampal network [5]. Theta oscillations have also been implicated in the encoding and retrieval of episodic and spatial memories [28, 41, 34, 11-13, 15-17] and disruption of them results in behavioural deficits [78]. Theta rhythm in hippocampus is paced by MS and diagonal band of Broca in the basal forebrain [66, 78], although several theta generators and theta dipoles seem to work independently in the hippocampus [5, 52].

Excitation and inhibition in hippocampus come in different flavours and supports different functions [20, 62]. Inhibition sculpts the activities of excitatory cells (GCs in DG and PCs in CA3 and CA1), allowing them to fire at particular temporal windows and phases with respect to external network oscillations [62, 51]. At least 25 different types of inhibitory interneurons have been identified in regions DG, CA3 and CA1 of the hippocampus [62, 63, 35, 21-23, 73]. These include axo-axonic cells (AACs), the perisomatic basket cells (BCs) and the dendritic bistratified cells (BSCs), ivy (IVY), neurogliaform (NGL), oriens lacunosum-moleculare (OLMs), molecular layer interneurons with axons in perforant-path termination zone (MOPP), hilar perforant path associated cells (HCs), hilar interneurons with axons in the commissural/associational pathway termination zone (HICAP), and the interneuron-selective cells (IS) in DG [63, 21-23, 9]. AACs innervate exclusively the initial axonal segment of the DG GCs and the CA3 and CA1 PCs, whereas BCs innervate their

cell bodies and proximal dendrites [63, 73]. CA1's BSCs and IVYs innervate the CA1 PC basal and oblique dendrites, whereas OLM and NGL cells target the apical dendritic tuft of CA3 and CA1 PCs aligned with the EC input [9, 62]. The DG HC cells target the apical dendrites of the DG GCs [73], whereas the MOPP cells feedforwardly inhibit the DG GCs. HICAP cells inhibit the proximal GC dendrites, near where MC axons terminate and provide feedback inhibition. IS cells inhibit exclusively other interneurons and modulate the excitability and synchrony of the network. Long range INs, such as the somatostatin- and mGluR1 α -positive neurons in stratum oriens project to the subiculum, other cortical areas and the medial septum, whereas somatostatin-negative ones and trilaminar cells project to the subiculum and other cortical areas but not to the septum [62-63, 35].

DG, CA3 and CA1 cells discharge at different phases of theta oscillations [63, 21-23, 9, 51, 62]. CA1 OLMs, BSCs, IVYs and PCs fire at the trough of theta recorded in the CA1 SP, whereas CA1 AACs, BCs and NGLs fire at the peak of theta recorded in the CA1 SP [63, 21-23]. CA3 AACs fire rhythmically around the peak of the theta oscillations recorded locally in CA3 [74], whereas CA3 BCs and PCs fire around the trough of the local CA3 theta with the PCs firing leading the BCs firing by few degrees [70]. CA3 OLMs, which are recurrently excited by the CA3 PCs should fire at the trough of CA3 theta right after the CA3 PCs. In addition to hippocampal cells, MS cell activities are theta modulated [67, 18, 3]. GABAergic MS neurons form two distinct populations exhibiting highly regular bursting activity that is coupled to either the trough or the peak of hippocampal theta waves [3].

In addition to theta oscillations, cells in the hippocampus fire at different phases of other rhythms, such as gamma (30-80Hz) and sharp wave-associated ripples (SWRs) (100-200Hz). Gamma oscillations constitute a basic clock cycle [25] and are embedded in theta oscillations [10]. SWRs occur during the offline replay and consolidation of previous experiences [62]. In CA1, during sharp wave ripple oscillations, BCs and BSCs strongly increase their discharge rates in phase with the ripple episode. In contrast, axo-axonic cells fire before the ripple episode, but pause their activities during and after it. OLM cells pause their firings during ripples. On the other hand, during theta oscillations, OLM cells, BSCs and PCs increase their firing rates at the troughs of the extracellular theta, whereas BCs and AACs fire at the peaks of it. During gamma oscillations, the firing rates of BCs, AACs and BSCs correlate with the extracellular gamma in different degrees, whereas OLM cells do not correlate at all with gamma oscillations.

Understanding the functional roles of these cells in encoding and retrieval of memories and rhythm generation currently poses a great challenge. Computational and mathematical models play an instrumental role in exploring such functions and facilitate the dissection of operations performed by the diverse interneurons. The aim of the next section is to provide a snapshot and a resumé of the current state-of-the-art of the ongoing research avenues concerning computational models of hippocampal microcircuits with particular emphasis on the functional roles of the various inhibitory interneurons in memory formation in the hippocampus in health and in disease.

COMPUTATIONAL MODELS OF HIPPOCAMPAL MICROCIRCUITS

Of paramount importance in memory research is the ability of any system to learn new things and acquire new memories, while at the same time not forgetting what it had previously learned. Such a memory system ought to be capable of determining which is new, requiring a distinct memory to be formed (pattern separation), and which details need to be recalled from incomplete or noisy information (pattern completion). In the next two sections I will review some of the most prominent microcircuit models of the hippocampus in relation to memory formation, storage and recall, in both health and disease.

Models of the healthy hippocampus

Sommer and Wennekers [60, 61] extended the original hippocampal CAM model [45, 76, 55] to investigate its memory capacity and robustness of efficient retrieval under varying memory load and type of external stimulation (tonic and pulsed). For learning they used the clipped synaptic modification rule of the Willshaw model [76]. Memory patterns were sequences of binary numbers (1 or 0). Each pattern was presented to a fixed number of cells in the network and each cell was active in more than one memory pattern. Inhibition worked as a global non-constant threshold. With tonic stimulation, the addressed memory was an attractor of the network dynamics. The memory was displayed rhythmically, coded by phase-locked bursts or regular spikes. The participating neurons had

rhythmic activity in the gamma-frequency range (30-80 Hz). If the input was switched from one memory to another, the network activity followed this change within one or two gamma cycles. With pulsed stimulation, memories were no longer attractors and they were retrieved within one or two gamma cycles. Burst of firing became relevant for coding and its occurrence was used for discriminating related processes from background activity.

Hunter and colleagues [33] compared and contrasted the performance of the Sommers and Wennekers model with previously published recall results of the Willshaw model [26, 27]. They tested how well the network can recall a pattern when there is full (100%) or partial (10%) connectivity or corruption due to noise and how the global inhibitory threshold could implement the winner-take-all (WTA) recall of a stored pattern. Biophysical implementations of three separate WTA recall methods were used: (1) Standard WTA implemented by intrinsic PC thresholding (increases in Na^+ density and membrane resistance) and global inhibition; (2) Normalized WTA implemented by localized inhibition proportional to the excitation a cell could receive, the range of EPSPs and the dendritic sums produced; and (3) amplified WTA via a non-linear increase of EPSP summation, so that the cells that reached a certain membrane potential increased their summed EPSP amplitude via a persistent Na^+ current. Recall was tested by tonically stimulating a subset of principal neurons in the network using an injected current of varying strength. Recall performance was tested by storing 50 random patterns, each consisting of 10 active cells, in the network and then using 5 of the 10 cells of a stored pattern as a recall cue. Recall quality with 10% connectivity was: (1) 61% in standard WTA, (2) 64% in normalized WTA and (3) 65% in amplified WTA.

Kunec et al. [41] advanced a detailed CA3 model of the hippocampus using biophysical representations of the major cell types including pyramidal cells (PCs) and two types of interneurons (BCs and OLMs) to dissect the operations performed by the various types of interneurons in and inputs to the network as well as investigate how variations in biophysically meaningful and experimentally measurable parameters affect the simulated encoding and retrieval. Inputs to the network came from MS, which paced the theta rhythm in the CA3 model into two half sub-cycles (one for storage and the other one for recall), and the EC (directly and via DG). Their model reproduced experimental results showing that the various cell types fire at a preferred phase relationship with respect to the underlying theta rhythm and to each other and offered distinct

functional roles of the various cells in storage and recall of memory patterns in CA3.

Drawing inspiration from the Kunec et al. [41] model and the experimental data of Klausberger and colleagues [38, 39], Cutsuridis and colleagues [11, 13, 12, 15, 16, 17] advanced a series of detailed biophysical models of the CA1 microcircuit of the hippocampus in order to investigate the memory capacity, recall performance, rate and phase coding properties and functional roles of various types of cells in region CA1 as a function of cue loading, presentation frequency and learning task. The Cutsuridis and colleagues modelling studies have been very influential in computational memory research and were the first to make a number of theoretical predictions regarding the roles of various types of inhibitory interneurons in the hippocampus in memory formation, which only recently have been experimentally validated:

- Theta modulated inhibition is what separates encoding and retrieval of memories in the hippocampus into two functionally independent half-cycles of the theta rhythm [11, 13]. This theoretical prediction has been recently experimentally verified [59].
- Theta modulated perisomatic inhibition plays an instrumental role in the encoding of memories in region CA1 of the hippocampus by allowing the generation of dendritic calcium spikes that promote synaptic LTP, while minimising cell output [11, 13]. This theoretical prediction has recently been experimentally verified [59].
- Theta modulated proximal dendritic inhibition in region CA1 of the hippocampus controls both cell output and suppresses dendritic calcium spikes, thus preventing LTP [11, 13]. This theoretical prediction has recently been experimentally verified [59].
- Theta modulated distal dendritic inhibition in region CA1 of the hippocampus removes interference from spurious memories during recall [11, 13].
- Intra- and extra-hippocampal inhibition provides the necessary environment for the maintenance of rate and phase coding properties of place cells in region CA1 of the hippocampus [16]. This theoretical prediction has been experimentally verified recently [36].

Recently, Nolan and colleagues [53] suggested that spike timing in the hippocampus is the mechanism capable of deciding when to learn a novel input pattern and when to recall by completing to a previously learned pattern, using a decision criterion based on patterns currently stored in the system. They advanced a novel computational model of the DG-CA3 microcircuit implicitly performing pattern-by-pattern novelty separation. The model incorporated spike timing dependent plasticity (STDP) as the mechanism to discriminate between known and unknown patterns by switching on and off learning. This pattern-by-pattern suppression ensured that even in unfamiliar situations, already known patterns are not relearned; whereas in familiar situations, unknown patterns could be learned. Simulation results demonstrated that (a) STDP in the EC-CA3 synapses provided a pattern completion ability without recurrent CA3 connections, (b) the race between activation of CA3 cells via EC-CA3 synapses and activation of the same cells via DG-CA3 synapses distinguished novel from known inputs, and (c) modulation of the EC-CA3 synapses adjusted the learned versus test input similarity required to evoke a direct CA3 response prior to any DG activity, thereby adjusting the pattern completion threshold.

Along the lines set by Nolan et al [53], Hummos and colleagues [32] advanced a more biologically realistic model of the DG-CA3 microcircuit that included principal cells and two of the most common interneurons, basket cells (BCs) and oriens lacunosum-moleculare (OLM) cells. Both inhibitory interneurons in the model were modulated by ACh according to experimental data [42]. The model suggested pattern separation and completion the DG-CA3 circuits produce instability through different dynamics, consequently requiring different mechanisms for their stabilization. Although the recurrent connections in CA3 promote runaway excitation, OLM inhibition and short-term depression at the recurrent connections are effective in preventing this instability, whereas BC inhibition by itself is not. Also, low ACh levels enhance CA3 recurrent connections leading to more sustained bursting in pyramidal cells, and short-term depression at these recurrent connections moderates this excitatory activity, whereas high ACh levels result in very long burst sizes that are optimally controlled by OLM inhibition.

Models of the diseased hippocampus

Menschik and Finkel [50] were the first to advance an Alzheimer's disease (AD) model of hippocampal CA3 region dynamics inspired by the Buzsaki "two-stage" memory model and the suggested role for interneurons [4, 7] and the Lisman and colleagues model on embedded gamma cycles within the theta rhythm [43, 44]. They used detailed biophysical representations of multi-compartmental models of pyramidal cells and two types of inhibitory interneurons: basket cells and chandelier cells to study the modulation and control of storage and recall dynamics in Alzheimer's disease by subcortical cholinergic and GABAergic input to the hippocampus. They showed that synchronization in the gamma frequency range can implement an attractor based auto-associative memory, where each new input pattern that arrives at the beginning of each theta cycle comprised of 5-10 embedded gamma cycles drives the network activity to converge over several gamma cycles to a stable attractor that represents the stored memory. Their results supported the hypothesis that spiking and bursting in CA3 pyramidal cells mediate separate behavioural functions and that cholinergic input regulates the transition between behavioural states associated with the online processing and recall of information. Cholinergic deprivation led to the slowing of gamma frequency, which reduced the number of "gamma cycles" within the theta rhythm available to reach the desired attractor state (i.e. memory loss and cognitive slowing seen in AD).

Inspired by the Cutsuridis and colleagues [13] modeling study, Bianchi et al. [2] investigated the conditions under which the properties of hippocampal CA1 pyramidal neuron altered by increasing CREB activity may contribute to improve memory storage and recall. With a set of patterns already stored in the network, they found that the pattern recall quality under AD-like conditions is significantly better when boosting CREB function with respect to control. Their results were robust even when synaptic damage due to AD progression increased, supporting the idea that the use of CREB-based therapies could provide a new approach to treat AD.

Yim and colleagues [79] extended an already well-established DG microcircuit model [58] into testing the hypothesis of whether the experimentally observed intrinsic scaling of GC activity serves as a

mechanism to maintain the pattern separation function of the DG network. They found that while increasing perirhinal path (PP; direct EC input to DG) strength degraded pattern separation only gradually, slight elevation of mossy fiber (MF) sprouting severely impaired pattern separation. When the DG network was hyper-excited, then the leaky GCs ameliorated pattern separation. In some sprouting cases with all-or-none seizure behavior, they observed pattern separation to be disabled with and without leaky GCs. When MF sprouting was mild (and PP strength was increased), then leaky GCs were particularly effective in restoring pattern separation performance.

CONCLUSIONS

I hope I made evident that large scale biophysical microcircuit models of associative memory are very important, because they allow us to run in-silico experiments of networks of neurons, while bypassing the technical difficulties of a real experiment, in order to answer questions and uncover mechanisms related to the interaction between the local microcircuit activity and global processing to achieve the desired overall processing functionality observed in learning and memory. Several practical issues such as parameter searching, network scaling, suitable simulation environments, and computational speed, memory and efficiency ought to be addressed in future large scale modeling studies of memory formation [8, 19, 24, 29, 71].

REFERENCES

1. Alonso, A., & Garcia-Austt, E. (1987). Neuronal sources of theta rhythm in the entorhinal cortex of the rat. I. Laminar distribution of theta field potentials. *Exp. Brain Res*, 67, 493–501
2. Bianchi, D., De Michele, P., Marchetti, C., Tirozzi, B., Cuomo, S., Marie, H., Migliore, M. (2014). Effects of increasing CREB-dependent transcription on the storage and recall processes in a hippocampal CA1 microcircuit. *Hippocampus*, 24(2), 165-77.
3. Borhegyi, Z., Varga, V., Szilagy, N., Fabo, D., Freund, T.F. (2004). Phase segregation of medial septal GABAergic neurons during hippocampal theta activity. *J Neurosci*, 24, 8470–8479

4. Buzsaki, G. (1989). Two-stage model of memory trace formation: a role for “noisy” brain states. *Neuroscience*, *31*, 551-70.
5. Buzsaki, G. (2002). Theta oscillations in the hippocampus. *Neuron*, *33*, 325-340
6. Buzsaki, G., Anastasiou, C., Koch, C. (2012). The origin of extracellular fields and currents: EEG, ECoG, LFP, and spikes. *Nature Rev Neurosci*, *13*(6), 407-20
7. Buzsaki, G., Chrobak, J. J. (1995). Temporal structure in spatially organized neuronal assemblies: A role for interneuronal networks. *Curr Opin Neurobiol*, *5*, 504-10
8. Carlson, K. D., Nageswaran, J. M., Dutt, N., Krichmar, J. L. (2014). An efficient automated parameter tuning framework for spiking neural networks. *Front Neurosci*, *8*, 10. doi: 10.3389/fnins.2014.00010.
9. Capogna, M. (2011). Neurogliaform cells and other interneurons of stratum lacunosum-moleculare gate entorhinal-hippocampal dialogue. *J Physiol*, *589*, 1875-1883
10. Colgin, L. L. (2015). Theta-gamma coupling in the entorhinal-hippocampal system. *Curr Opin Neurobiol*. *31*, 45-50. doi: 10.1016/j.conb.2014.08.001
11. Cutsuridis, V., Cobb, S., Graham, B. P. (2008). Encoding and Retrieval in a CA1 Microcircuit Model of the Hippocampus. In V. Kurkova et al. (Eds.), *Lecture Notes in Computer Science* (LNCS 5164, pp. 238-247). Berlin Heidelberg: Springer-Verlag.
12. Cutsuridis, V., Wenneckers, T. (2009). Hippocampus, microcircuits and associative memory. *Neural Networks*, *22*(8), 1120-8
13. Cutsuridis, V., Graham, B. P., Cobb, S. (2010a). Encoding and retrieval in the hippocampal CA1 microcircuit model. *Hippocampus*, *20*(3), 423-446
14. Cutsuridis, V., Graham, B. P., Cobb, S., Vida, I. (2010b). *Hippocampal Microcircuits: A Computational Modeler's Resource Book*. New York: Springer.
15. Cutsuridis, V., Graham, B. P., Cobb, S., Hasselmo, M. E. (2011). Bio-inspired models of memory capacity, recall performance and theta phase precession. *Proceedings of International Joint Conference on Neural Networks*, San Jose, California, USA, July 31- August 5, pp. 3141-48
16. Cutsuridis, V., Hasselmo, M. (2012). GABAergic modulation of gating, timing and theta phase precession of hippocampal neuronal activity during theta oscillations. *Hippocampus*, *22*, 1597-1621
17. Cutsuridis, V., Poirazi, P. (2015). A computational study on how theta modulated inhibition can account for the long temporal delays in the entorhinal-hippocampal loop. *Neurobiology of Learning and Memory*, *120*, 69-83
18. Dragoi, G., Carpi, D., Recce, M., Csicsvari, J., Buzsaki, G. (1999). Interactions between hippocampus and medial septum during sharp waves and theta oscillation in the behaving rat. *J Neurosci*. *19*, 6191-6199
19. Djurfeldt, M., Ekeberg, O., & Lansner, A. (2008). Large-scale modeling—a tool for conquering the complexity of the brain. *Front. Neuroinformatics* *2*,1. doi: 10.3389/neuro.11.001.2008
20. Freund, T.F., Buzsaki, G. (1996). Interneurons of the hippocampus. *Hippocampus*, *6*, 347-470
21. Fuentealba, P., Begum, R., Capogna, M., Jinno, S., Marton, L. F., Csicsvari, J., Thomson, A., Somogyi, P., & Klausberger, T. (2008a). Ivy cells: A population of nitric-oxide-producing, slow-spiking GABAergic neurons and their involvement in hippocampal network activity. *Neuron*, *57*, 917-929
22. Fuentealba, P., Klausberger, T., Karayannis, T., Suen, W. Y., Huck, J., Tomioka, R., Rockland, K., Capogna, M., Studer, M., Morales, M., Somogyi, P. (2010). Expression of COUP-TFII nuclear receptor in restricted GABAergic neuronal populations in the adult rat hippocampus. *J. Neurosci*, *30*, 1595-1609
23. Fuentealba, P., Tomioka, R., Dalezios, Y., Marton, L. F., Studer, M., Rockland, K., Klausberger, T., Somogyi, P. (2008b). Rhythmically active enkephalin- expressing GABAergic cells in the CA1 area of the hippocampus project to the subiculum and preferentially innervate interneurons. *J Neurosci*, *28*, 10017-10022
24. Gleeson, P., Silver, R.A., Steuber, V. (2010). Computer simulation environments. In V. Cutsuridis, B. P. Graham, S. Cobb, I. Vida (Eds.), *Hippocampal microcircuits: A computational modeller's resource book*. New York: Springer
25. Graham, B. P. (2003). Dynamics of storage and recall in hippocampal associative memory networks. In P. Erdi, A. Esposito, M. Marinaro, S. Scarpetta (Eds.), *Computational Neuroscience: Cortical Dynamics* (1st edition, pp. 1-23). Berlin Heidelberg: Springer-Verlag
26. Graham, B., & Willshaw, D. (1995). Improving recall from an associative memory. *Biological Cybernetics*, *72*, 337-346.
27. Graham, B., & Willshaw, D. (1997). Capacity and information efficiency of the associative net. *Network*, *8*, 35-54

28. Hasselmo, M. (2005). What is the function of hippocampal theta rhythm? – Linking behavioural data to phasic properties of field potential and unit recording data. *Hippocampus*, *15*, 936-949
29. Hasselmo, M. E., Kapur, A. (2000). Modelling of large networks. In E. De Schutter (Ed.), *Computational neuroscience: realistic modelling for experimentalists* (1st edition, pp. 289-315). CRC Press
30. Hasselmo, M. E., Schnell, E. (1994). Laminar selectivity of the cholinergic suppression of synaptic transmission in rat hippocampal region CA1: computational modeling and brain slice physiology. *J. Neurosci*, *14*, 3898-3914
31. Hasselmo, M. E., & Wyble, B. P. (1997). Free recall and recognition in a network model of the hippocampus: simulating effects of scopolamine on human memory function. *Behav Brain Res*, *89*, 1-34
32. Hummos, A., Franklin, C. C., Nair, S. S. (2014). Intrinsic mechanisms stabilize encoding and retrieval circuits differentially in a hippocampal network model. *Hippocampus*, *24*(12), 1430-48. doi: 10.1002/hipo.22324
33. Hunter, R., Cobb, S., & Graham, B. P. (2008). Improving associative memory in a network of spiking neurons. In V. Kurkova et al. (Eds.), *Lecture Notes in Computer Science* (LNCS 5164, pp. 636_645). Berlin Heidelberg: Springer-Verlag.
34. Jensen, O., & Lisman, J. E. (2005). Hippocampal sequence-encoding driven by a cortical multi-item working memory buffer. *TINS*, *28*, 67–72
35. Jinno, S., Klausberger, T., Marton, L. F., Dalezios, Y., et al. (2007). Neuronal diversity in GABAergic long-range projections from the hippocampus. *J. Neurosci*. *27*(33), 8790-8804
36. Kaifosh, P., Lovett-Barron, M., Turi, G. F., Reardon, T.R., Losonczy, A. (2013). Septo-hippocampal GABAergic signaling across multiple modalities in awake mice. *Nat Neurosci*, *16*(9), 1182-4. doi: 10.1038/nn.3482.
37. Klausberger, T., Somogyi, P. (2008). Neuronal diversity and temporal dynamics: the unity of hippocampal circuit operations. *Science*, *321*, 53-57
38. Klausberger, T., Magill, P. J., Marton, L. F., David, J., Roberts, B., Cobden, P. M., Buzsaki, G., Somogyi, P. (2003). Brain-state- and cell-type-specific firing of hippocampal interneurons in vivo. *Nature*, *421*, 844-848
39. Klausberger, T., Marton, L. F., Baude, A., Roberts, J. D., Magill, P. J., Somogyi, P. (2004). Spike timing of dendrite-targeting bistratified cells during hippocampal network oscillations in vivo. *Nat Neurosci*. *7*(1), 41-7.
40. Kohonen, T. (1978). *Associative memory*. Springer-Verlag.
41. Kunec, S., Hasselmo, M. E., Kopell, N. (2005). Encoding and retrieval in the CA3 region of the hippocampus: a model of theta-phase separation. *J Neurophysiol*, *94*(1), 70-82.
42. Lawrence, J. J. (2008). Cholinergic control of GABA release: emerging parallels between neocortex and hippocampus. *TINS*, *31*(7), 317-27. doi: 10.1016/j.tins.2008.03.008
43. Lisman, J. (2005). The theta/gamma discrete phase code occurring during the hippocampal phase precession may be a more general brain coding scheme. *Hippocampus*, *15*, 913-922.
44. Lisman, J., Idiart, M.A. (1995). Storage of 7±2 short-term memories in oscillatory subcycles. *Science*, *267*, 1512-5.
45. Marr, D. (1971). A simple theory of archicortex. *Philos Trans R Soc Lond B Biol Sci.*, *262*(841), 23-81.
46. Marr, D. (1969). A theory of cerebellar cortex. *J. Physiol. (London)*, *202*, 437-70
47. McClelland, J. L., McNaughton, B. L., O'Reilly, R. C. (1995). Why there are complementary learning systems in the hippocampus and neocortex: insights from the successes and failures of connectionist models of learning and memory. *Psychol Rev*, *102*, 419-457.
48. McCulloch, W. S., Pitts, W. (1943). A logical calculus of the ideas immanent of the nervous activity. *Bull Math Biophys*, *5*, 115-133
49. McNaughton, B. L., Morris, R. G. M. (1987). Hippocampal synaptic enhancement and information storage within a distributed memory system. *TINS*, *10*(10), 408-415
50. Menschik, E. D., Finkel, L. H. (1998). Neuromodulatory control of hippocampal function: towards a model of Alzheimer's disease. *Artificial Intelligence in Medicine*, *13*, 99-121
51. Mizuseki, K., Sirota, A., Pastalkova, E., Buzsaki, G. (2009). Theta oscillations provide temporal windows for local circuit computation in the entorhinal-hippocampal loop. *Neuron*, *64*, 267-280
52. Montgomery, S. M., Betancur, M. I., Buzsaki, G. (2009). Behavior-dependent coordination of multiple theta dipoles in the hippocampus. *J. Neurosci*, *29*, 1381–1394.
53. Nolan, C. R., Wyeth, G., Milford, M., Wiles, J. (2011). The race to learn: spike timing and STDP can coordinate learning and recall in CA3. *Hippocampus*, *21*(6), 647-60. doi: 10.1002/hipo.20777

54. Palm, G. (1982). *Neural assemblies: an alternative approach to artificial intelligence*. Berlin: Springer-Verlag
55. Palm, G. (1980). On associative memories. *Biol. Cybern.* 36, 9-31
56. Palm, G. (1991). Memory capacities of local rules for synaptic modification. *Concepts in Neuroscience*, 2, 97-128
57. Palm, G, Sommer, F. (1996). Associative data storage and retrieval in neural networks. In E. Domany, J. van Hemmen, K. Schulten (Eds.), *Models of neural networks* (volume II, pp. 79-118). New York: Springer-Verlag
58. Santhakumar, V., Aradi, I., Soltesz, I. (2005). Role of mossy fiber sprouting and mossy cell loss in hyperexcitability: A network model of the dentate gyrus incorporating cell types and axonal topography. *J Neurophysiol*, 93, 437-453
59. Siegle, J.H., Wilson, M.A. (2014). Enhancement of encoding and retrieval functions through theta phase-specific manipulation of hippocampus. *eLife*, e03061
60. Sommer, F.T., Wennekers, T. (2000). Modelling studies on the computational function of fast temporal structure in cortical circuit activity. *J Physiol (Paris)*, 94, 473-488
61. Sommer, F. T., Wennekers, T. (2001). Associative memory in networks of spiking neurons. *Neural Networks*, 14, 825-834
62. Somogyi, P., Katona, L., Klausberger, T., Lasztocki, B., Viney, T. J. (2014). Temporal redistribution of inhibition over neuronal subcellular domains underlies state-dependent rhythmic change of excitability in the hippocampus. *Phil. Trans R Soc B*, 369, 20120518
63. Somogyi, P., Klausberger, T. (2005). Defined types of cortical interneurons structure space and spike timing in the hippocampus. *J Physiol*, 562, 9-26
64. Steinbuch, K. (1961). Non-digital learning matrices as preceptors. *Kybernetik*, 1, 117-124
65. Stella, F., Treves, A. (2011). Associative memory storage and retrieval: involvement of theta oscillations in hippocampal information processing. *Neural Plasticity*, 683961.
66. Stewart, M., Fox, S. (1990). Do septal neurons pace the hippocampal theta rhythm? *TINS*, 13, 163-169
67. Stumpf, C., Petsche, H., Gogolak, G. (1962). The significance of the rabbit's septum as a relay station between the midbrain and the hippocampus. II. The differential influence of drugs upon both the septal cell firing pattern and the hippocampus theta activity. *Electroencephalogr Clin Neurophysiol*, 14, 212-9.
68. Suzuki, W.A. (2007). *Ann N Y Acad Sci*, 1097, 1-11
69. Treves, A., Rolls, E. (1992). Computational constraints suggest the need for two distinct input systems to the hippocampal CA3 network. *Hippocampus*, 2, 189-200
70. Tukker, J. J., Lasztocki, B., Katona, L., Roberts, J. D. B., Pissadaki, E. K., Dalezios, Y., Marton, L., Zhang, L., Klausberger, T., Somogyi, P. (2013). Distinct dendritic arborisation and in vivo firing patterns of parvalbumin-expressing basket cells in the hippocampal area CA3. *J Neurosci*, 33(16), 6809-6825
71. Van Geit, W., de Schutter, E., Archard, P. (2008). Automated neuron model optimization techniques: a review. *Biol. Cybern*, 99, 241-251
72. Vanderwolf, C. H. (1969). Hippocampal electrical activity and voluntary movement in the rat. *Electroencephalogr. Clin Neurophysiol*, 26, 407-418
73. Vida, I. (2010). Morphology of hippocampal neurons. In V Cutsuridis, B.P. Graham, S. Cobb, I. Vida (Eds.), *Hippocampal Microcircuits: A Computational Modeler's Resource Book* (1st edition, pp 27-67). New York: Springer
74. Viney, T. J., Lasztocki, B., Katona, L., Crump, M.G., Tukker, J. J., Klausberger, T., Somogyi, P. (2013). Network state-dependent inhibition of identified hippocampal CA3 axo-axonic cells in vivo. *Nat Neurosci*, 16(12), 1802-1811
75. Vinogradova, O. (2001). Hippocampus as a comparator: Role of the two input and two output systems of the hippocampus in selection and registration of information. *Hippocampus*, 11, 578-598
76. Willshaw, D., Buneman, O., Longuet-Higgins, H. (1969). Non-holographic associative memory. *Nature*, 222, 960-962
77. Wilson, M.A., McNaughton, BL. (1993). Dynamics of the hippocampal ensemble code for space. *Science*, 261(5124), 1055-8
78. Winson, J. (1978). Loss of hippocampal theta rhythm results in spatial memory deficit in the rat. *Science*, 210, 160-163
79. Yim, M.Y., Hanuschkin, A., Wolfart, J. (2015). Intrinsic rescaling of granule cells restores pattern separation ability of a dentate gyrus network model during epileptic hyperexcitability. *Hippocampus*, 25(3), 297-308. doi: 10.1002/hipo.22373